PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/00, 9/12	A1	(11) International Publication Number:	WO 96/18384
		(43) International Publication Date:	20 June 1996 (20.06.96)
(21) International Application Number: PCT/EP9 (22) International Filing Date: 8 December 1995 (0 (30) Priority Data: 9425160.0 10 December 1994 (10.12.94) (71) Applicant (for all designated States except US): GROUP LIMITED [GB/GB]; Glaxo Wellcome Berkeley Avenue, Greenford, Middlesex UB6 0NN (72) Inventors; and	8.12.95) GI GLAXO House	CH, CN, CZ, DE, DK, EE, ES, KE, KG, KP, KR, KZ, LK, LR MK, MN, MW, MX, NO, NZ, SG, SI, SK, TJ, TM, TT, UA, U patent (AT, BE, CH, DE, DK, ES MC, NL, PT, SE), OAPI patent GA, GN, ML, MR, NE, SN, TD LS, MW, SD, SZ, UG).	FI, GB, GE, HU, IS, JP, LT, LU, LV, MD, MG, PL, PT, RO, RU, SD, SE G, US, UZ, VN, European G, FR, GB, GR, EE, IT, LU, (BF, BJ, CF, CG, CI, CM,
(75) Inventors/Applicants (for US only): SAPSFORD, [GB/GB]; Glaxo Research and Development, Part Ware, Hertfordshire SG12 0DP (GB). SAVAGE, / Patrick [GB/GB]; Glaxo Research and Development Road, Ware, Hertfordshire SG12 0DP (GB).	k Road, Andrew,		
(74) Agent: DAWSON, Hugh, B.; Glaxo Wellcome plc, Wellcome House, Berkeley Avenue, Greenford, Mi UB6 0NN (GB).			

(57) Abstract

This invention relates to aerosol formulations of use the administration of medicaments by inhalation in particular a pharmaceutical aerosol formulation which comprises (a) 1,1,1,2-tetrafluoroethane or 1,1,1,2,3,3,3-heptafluoro-n-propane or mixtures thereof as propellant, (b) 1,1,2,2,3-pentafluoropropane as co-propellant, and (c) particulate medicament. A method of treating respiratory disorders which comprises administration by inhalation of an effective amount of a pharmaceutical aerosol formulation as defined is also described.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

					•
AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	CE	Georgia .	MW	Melawi
88	Barbados	GN	Guinea	NE	Niger
BE	Belgium	. GR	Grecce	NL	Netherlands
BF	Burkina Feso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL.	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CG	Congo		of Kores	SE	Sweden
CH .	Switzerland .	. KR	Republic of Kores	SI.	Slovenia
a	Côse d'Ivoire	KZ.	Kazakhutan	SK	Slovakia
CM	Cameroon	ш	Liechtenstzin	5N	Senegal
CN	China	LK	Sri Lanka	170	Ched
CS	Czechoslovakia	LU	Luxembourg	TG	Tago
CZ	Czech Republic	LV	Larvia	TJ	Tajikistan
30	Germany	MC	Monaco	π	Trinided and Tobego
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ZS	Spein	MG	Madagascar	US	United States of America
n	Finland	ML	Mali	UZ	Usbekistan
J.R	France	MN	Mongolia	VN	Viet Nam
5.	Cohen			•••	

PROPELLANT MIXTURE FOR AEROSOL FORMULATION

This invention relates to aerosol formulations of use in the administration of medicaments by inhalation.

5

The use of aerosols to administer medicaments has been known for several decades. Such aerosols generally comprise the medicament, one or more chlorofluorocarbon propellants and either a surfactant or a solvent, such as ethanol.

10

The most commonly used aerosol propellants for medicaments have been CCl₃F (propellant 11) in admixture with CCl₂F₂ (propellant 12) and CF₂Cl.CF₂Cl (propellant 114). However these propellants are now believed to provoke the degradation of stratospheric ozone and there is thus a need to provide aerosol formulations for medicaments which employ so called "ozone-friendly" propellants.

20

15

A class of propellants which are believed to have minimal ozone-depleting effects in comparison to conventional chlorofluorocarbons comprise hydrogen-containing chlorofluorocarbons and fluorocarbons and a number of medicinal aerosol formulations using such propellant systems have been disclosed in, for example, EP 0372777, WO91/04011, W091/11173, W091/11495, W091/14422, W092/00061, W092/00062 and W092/00107.

25

30

These applications are all concerned with the preparation of pressurised aerosols for the administration of medicaments and seek to overcome the problems associated with the use of the new class of propellants, in particular the problems of stability associated with the pharmaceutical formulations prepared. These applications all propose the addition of a wide range of adjuvants such as alcohols, alkanes, dimethyl ether, surfactants (including fluorinated and non-fluorinated surfactants, carboxylic acids, polyethoxylates etc) and even conventional chlorofluorocarbon propellants in small amounts to minimise potential ozone damage.

Surprisingly, we have now found that mixtures of a non ozone-depleting propellant and a specific fluorinated hydrocarbon may be employed as propellant systems suitable for use in pharmaceutical aerosol compositions.

- There is thus provided in one aspect of the invention an aerosol formulation comprising:
 - (a) 1,1,1,2-tetrafluoroethane (CF₃CH₂F), 1,1,1,2,3,3,3-heptafluoro-n-propane (CF₃CHFCF₃) or mixtures thereof as propellant;
 - (b) 1,1,2,2,3-pentafluoropropane as co-propellant; and
- 10 (c) particulate medicament.

Generally, the ratio of propellant: co-propellant is in the range of about 30: 70 to about 95: 5, preferably 50: 50 to 90: 10 by weight, especially 50: 50 to 80: 20, for example 75: 25 (w/w).

15

20

25

30

35

Medicaments which may be administered in aerosol formulations according to the invention include any drugs useful in inhalation therapy which may be presented in a form which is substantially completely insoluble in the selected propellant system. Appropriate medicaments may thus be selected from, for example, analgesics, e.g. codeine, dihydromorphine, ergotamine, fentanyl or morphine; anginal preparations, e.g. dilitiazem; antiallergics, e.g. cromolyn, cromogylcate or nedocromil; antibiotics, e.g. cephalosporins, penicillins, antihistamines, tetracyclines; streptomycin, sulphonamides beclomethasone, flunisolide, anti-inflammatories, methapyrilene; e.g. fluticasone, tipredane, budesonide, triamcinolone acetonide; antitussives, e.g. noscapine: bronchodilators, e.g. ephedrine, epinephrine, fenoterol, formoterol, metaproterenol. isoprenaline. isoproterenol. phenylephrine, phenylpropanolamine, pirbuterol, repoterol, rimiterol, salbutamol, salmeterol, (-)-4-amino-3,5-dichloro-α-{[[6-{2-(2terbutaline pyridinyl)ethoxy]hexyl]amino]methyl]benzenemethanol; diuretics, e.g. amiloride; anticholinergics e.g. ipratropium bromide; hormones, e.g. hydrocortisone or prednisolone; and therapeutic proteins and peptides, e.g. glucagon or insulin. It will be clear to a person skilled in the art that, where appropriate, the medicaments will be used in the form of salts (e.g. as alkali metal or amine salts or as acid addition salts) or as esters (e.g. lower alkyl

10

15

25

30

35

esters) or as solvates (eg hydrates) to optimise the activity and/or stability of the medicament and/or to minimise the solubility of the medicament in the propellant.

Particularly preferred medicaments for administration using aerosol formulations in accordance with the invention include bronchodilators and anti-inflammatory steroids of use in the treatment of asthma by inhalation therapy, for example salbutamol (e.g. as the sulphate), salmeterol (e.g. as the hydroxynaphthoate known as salmeterol xinafoate), becomethasone dipropionate or a solvate thereof, fluticasone propionate or (-)-4-amino-3,5-dichloro-α-[[[6-[2-(2-yyridinyl)ethoxy]hexyl]amino]methyl] benzenemethanol.

The particle size of the particulate medicament should be such as to permit inhalation of substantially all of the medicament into the lungs upon administration of the aerosol formulation and will thus desirably be less than 20 microns, preferably in the range 1 to 10 microns, e.g. 1 to 5 microns. The particle size of the medicament may be reduced by conventional means, for example by milling or micronisation.

The final aerosol formulation desirably contains 0.005-10% w/w, preferably 0.005-5% w/w, especially 0.01-1.0% w/w, of medicament relative to the total weight of the formulation.

It is desirable that the formulations of the invention contain no components which may provoke the degradation of stratospheric ozone. In particular it is desirable that the formulations are substantially free of chlorofluorocarbons such as CCl₃F, CCl₂F₂ and CF₃CCl₃. As used herein "substantially free" means less than 1% w/w based upon the propellant system, in particular less than 0.5%, for example 0.1% or less.

The propellant may optionally contain an adjuvant having a higher polarity and/or a higher boiling point than the propellant. Polar adjuvants which may be used include (e.g. C₂₋₆) aliphatic alcohols and polyols such as ethanol, isopropanol and propylene glycol, preferably ethanol. In general only small quantities of polar adjuvants (e.g. 0.05 - 3.0% w/w) may be required to improve

the stability of the dispersion - the use of quantities in excess of 5% w/w may tend to dissolve the medicament. Formulations in accordance with the invention may preferably contain less than 1% w/w, e.g. about 0.1% w/w, of polar However, the formulations of the invention are preferably substantially free of polar adjuvants, especially ethanol. Suitable volatile adjuvants include saturated hydrocarbons such as propane, n-butane, isobutane, pentane and isopentane and alkyl ethers such as dimethyl ether. In general, up to 50% w/w of the propellant may comprise a volatile edjuvant, for example 1 to 30% w/w of a volatile saturated C1-6 hydrocarbon.

10

15

5

Optionally, the aerosol formulations according to the invention may further comprise one or more surfactants. The surfactants must be physiologically acceptable upon administration by inhalation. Within this category are included surfactants such as oleic acid, sorbitan trioleate (Span R 85), sorbitan monooleate, sorbitan monolaurate, polyoxyethylene (20) sorbitan monolaurate, lecithin, oleyl natural monooleate, (20) sorbitan polyoxyethylene polyoxyethylene (2) ether, stearyl polyoxyethylene (2) ether, lauryl polyoxyethylene (4) ether, block copolymers of oxyethylene and oxypropylene, synthetic lecithin, diethylene glycol dioleate, tetrahydrofurfuryl oleate, ethyl oleate, isopropyl myristate, glyceryl monooleate, glyceryl monostearate, glyceryl monoricinoleate, cetyl alcohol, steanyl alcohol, polyethylene glycol 400, cetyl pyridinium chloride, benzalkonium chloride, olive oil, glyceryl monolaurate, com oil, cotton seed oil and sunflower seed oil. Preferred surfactants are lecithin, oleic acid and sorbitan trioleate.

25

20

An alternative class of surfactants are described in EP 0478686, especially surfactants of formula (I)

native class of surfactants are described in EP 0478686, esents of formula (I)

$$C_{n}F_{2n-1}(CH_{2})_{m}-C-O-CH_{2}$$

$$O \qquad (I)$$

$$C_{n}F_{2n-1}(CH_{2})_{m}-C-O-CH$$

$$C_{n}F_{2n-1}(CH_{2})_{m}-C-O-CH$$

$$C_{n}F_{2n-1}(CH_{2})_{m}-C-O-CH$$

$$CH_{2}-O-CH_{2}CH_{2}CH_{2}N-R^{2}$$

$$CH_{2}-O-CH_{2}CH_{2}CH_{2}N-R^{2}$$

wherein n is an integer of 1 to 18, especially 2 to 12; m is an integer of 0 to 17, especially 0 to 11; and R¹, R² and R³ are each independently a hydrogen atom or a C₁₋₄alkyl group.

5

Particularly preferred surfactants of formula (I) are the fluorinated phosphatidylcholines wherein R^1 , R^2 and R^3 each represent methyl, n is an integer of 4 to 8, especially 4 or 6, and m is an integer of 4 to 10, especially 4 or 6.

10

If desired, the surfactant may be incorporated into the aerosol formulation in the form of a surface coating on the particulate medicament. In this case, the use of substantially non-ionic surfactants which have reasonable solubility in substantially non-polar solvents is frequently advantageous since it facilitates coating of the medicament particles using solutions of surfactant in non-polar solvents in which the medicament has limited or minimal solubility.

20

15

The amount of surfactant employed in coating the particulate medicament is desirably in the range 0.1 to 10% w/w, preferably 1 to 10% w/w, relative to the medicament. Where the surfactant is present as a surface coating, the amount may advantageously be chosen such that a substantially monomolecular coating of surfactant is formed. However, it is preferable that the formulations of the invention are substantially free of surfactants, i.e. contain less than an effective stabilising amount of a surfactant such as less than 0.0001% by weight of medicament.

25

30

The formulations of the invention may be prepared by dispersal of the medicament in the selected propellant and/or co-propellant in an appropriate container, e.g. with the aid of sonication. Preferably the particulate medicament is suspended in co-propellant and filled into a suitable container. The valve of the container is then sealed into place and the propellant introduced by pressure filling through the valve in the conventional manner. Surprisingly, the aerosol formulations according to the invention have been found to be easily redispersed by mild agitation to provide suspensions with excellent delivery

10

15

20

25

characteristics suitable for use in pressurised inhalers, even after prolonged storage.

The chemical and physical stability and the pharmaceutical acceptability of the aerosol formulations according to the invention may be determined by techniques well known to those skilled in the art. Thus, for example, the chemical stability of the components may be determined by HPLC assay, for example, after prolonged storage of the product. Physical stability data may be gained from other conventional analytical techniques such as, for example, by leak testing, by valve delivery assay (average shot weights per actuation), by dose reproducibility assay (active ingredient per actuation) and spray distribution analysis.

The formulations according to the invention may be filled into canisters suitable Canisters generally for delivering pharmaceutical aerosol formulations. comprise a container capable of withstanding the vapour pressure of the propellant used such as a plastic or plastic-coated glass bottle or preferably a metal can, for example an aluminium can which may optionally be anodised, lacquer-coated and/or plastic-coated, which container is closed with a metering valve. The metering valves are designed to deliver a metered amount of the formulation per actuation and incorporate a gasket to prevent leakage of The gasket may comprise any suitable propellant through the valve. elastomeric material such as for example low density polyethylene, chlorobutyl, black and white butadiene-acrylonitrile rubbers, butyl rubber and neoprene. Suitable valves are commercially available from manufacturers well known in the aerosol industry, for example, from Valois, France (e.g. DF10, DF30, DF60), Bespak plc, UK (e.g. BK300, BK356) and 3M-Neotechnic Ltd, UK (e.g. SpraymiserTM).

Conventional bulk manufacturing methods and machinery well known to those skilled in the art of pharmaceutical aerosol manufacture may be employed for the preparation of large scale batches for the commercial production of filled canisters. Thus, for example, in one bulk manufacturing method the particulate medicament is first suspended in the co-propellant. The drug suspension is then filled into the empty canisters, valves crimped on and then propellant is

pressure filled into the canisters through the valves in conventional manner. Typically, in batches prepared for pharmaceutical use, each filled canister is check-weighed, coded with a batch number and packed into a tray for storage before release testing.

5

Each filled canister is conveniently fitted into a suitable channelling device prior to use to form a metered dose inhaler for administration of the medicament into the lungs or nasal cavity of a patient. Suitable channelling devices comprise for example a valve actuator and a cylindrical or cone-like passage through which medicament may be delivered from the filled canister via the metering valve to the nose or mouth of a patient e.g. a mouthpiece actuator. Metered dose inhalers are designed to deliver a fixed unit dosage of medicament per actuation or "puff", for example in the range of 10 to 5000 microgram medicament per puff.

15

20

25

10

Administration of medicament may be indicated for the treatment of mild, moderate or severe acute or chronic symptoms or for prophylactic treatment. It will be appreciated that the precise dose administered will depend on the age and condition of the patient, the particular particulate medicament used and the frequency of administration and will ultimately be at the discretion of the attendant physician. When combinations of medicaments are employed the dose of each component of the combination will in general be that employed for each component when used alone. Typically, administration may be one or more times, for example from 1 to 8 times per day, giving for example 1,2,3 or 4 puffs each time.

The following non-limitative Examples serve to illustrate the invention.

Example 1

30

35

Micronised salmeterol xinafoate (hydroxynaphthoate, 8.7mg) was weighed into a clean, dry glass bottle together with 1,1,2,2,3-pentafluoropropane (1.3g, 1ml). The bottle was sealed by crimping a valve in place. Propellant 1,1,1,2-tetrafluoroethane (20.7g, 17ml) was added, under pressure, through the valve. The resultant inhaler delivers 25µg of salmeterol xinafoate (hydroxynaphthoate)

per actuation (200 75mg actuations per bottle). The ratio of propellant (CF_3CH_2F) to co-propellant $(CHF_2CF_2CH_2F)$ was 17:1 (v/v).

Example 2

5

Micronised salmeterol xinafoate (hydroxynaphthoate, 8.7mg) was weighed into a clean, dry glass bottle together with 1,1,2,2,3-pentafluoropropane (5.2g, 4ml). The bottle was sealed by crimping a valve in place. Propellant 1,1,1,2-tetrafluoroethane (17.1g, 14ml) was added, under pressure, through the valve. The resultant inhaler delivers 25 μ g of salmeterol xinafoate (hydroxynaphthoate) per actuation (200 75mg actuations per bottle). The ratio of propellant (CF₃CH₂F) to ∞ -propellant (CHF₂CF₂CH₂F) was 14 : 4 (v/v).

Example 3

15

20

10

Micronised salmeterol xinafoate (hydroxynaphthoate, 4mg) was weighed into a clean, dry aluminium aerosol canister (8ml) together with co-propellant 1,1,2,3-pentafluoropropane (2g). The canister was sealed by crimping a valve in place and propellant 1,1,1,2-tetrafluoroethane (10g) was added, under pressure, through the valve. The resultant inhaler delivers 25µg of salmeterol hydroxynaphthoate per actuation (120 75mg actuations per can).

Examples 4 to 7

25

Inhalers were prepared as described in Example 3 containing propellant (CF_3CH_2F) to ∞ -propellant $(CHF_2CF_2CH_2F)$ in the ratios of 9:3, 8:4, 7:5 and 6:6 (w/w) (Examples 4, 5, 6 and 7 respectively).

Example 8

30

35

Micronised salbutamol (base) (24mg) is homogenised with the aid of sonication in a solution of oleic acid (2.4mg) in the co-propellant 1,1,2,2,3-pentafluoropropane (4.7g) and filled into a clean, dry aluminium aerosol canister. The canister is sealed by crimping a valve in place. Propellant 1,1,1,2-tetrafluoroethane (14.1g) is added, under pressure, through the valve.

The resultant inhaler delivers 100 microgram salbutamol per 75mg actuation. The ratio of propellant (CF₃CH₂F) to co-propellant (CHF₂CF₂CH₂F) was 75 : 25 (w/w).

5 Examples 9 to 11

Inhalers are prepared as described in Example 8 containing propellant (CF_3CH_2F) and co-propellant $(CHF_2CF_2CH_2F)$ in the ratios 70 : 30, 50 : 50 and 95 : 5 (w/w) (Examples 9, 10 and 11 respectively).

Example 12

10

15

20

Micronised salbutamol (base) (24mg) is homogenised with the aid of sonication in a solution of oleic acid (2.4mg) in the co-propellant 1,1,2,2,3-pentafluoropropane (5.3g) and filled into a clean, dry aluminium aerosol canister. The canister is sealed by crimping a valve in place. Propellant 1,1,1,2,3,3,3-heptafluoro-n-propane (15.9g) is added, under pressure, through the valve. The resultant inhaler delivers 100 microgram salbutamol per 75mg actuation. The ratio of propellant (CF3CHFCF3) to co-propellant (CHF2CF2CH2F) was 75: 25 (w/w).

Examples 13 to 15

Inhalers are prepared as described in Example 12 containing propellant (CF₃CHFCF₃) to co-propellant (CHF₂CF₂CH₂F) in the ratios 70 : 30, 50 : 50 and 95 : 5 (w/w) (Examples 13, 14 and 15 respectively).

Example 16

Micronised fluticasone propionate (4mg) is weighed into a clean, dry aluminium aerosol canister (8ml) together with co-propellant 1,1,2,2,3-pentafluoropropane (3.1g). The canister is sealed by crimping a valve in place. Propellant 1,1,1,2-tetrafluoroethane (9.3g) is added, under pressure, through the valve. The resultant inhaler delivers 25μg of fluticasone propionate per actuation (120)

75mg actuations per can). The ratio of propellant (CF₃CH₂F) to ∞ -propellant (CHF₂CF₂CH₂F) is 9:3w/w.

Example 17

5

10

Micronised salmeterol xinafoate (hydroxynaphthoate, 4mg) and micronised fluticasone propionate (8mg) are weighed into a clean, dry aluminium aerosol canister (8ml) together with co-propellant 1,1,2,2,3-pentafluoropropane (3.1g). The canister is sealed by crimping a valve in place. Propellant 1,1,1,2-tetrafluoroethane (9.3g) is added, under pressure, through the valve. The resultant inhaler delivers 25μg salmeterol xinafoate (hydroxynaphthoate) and 50μg fluticasone propionate per actuation (120 75mg actuations per can). The ratio of propellant (CF₃CH₂F) to co-propellant (CHF₂CF₂CH₂F) is 9:3w/w.

CLAIMS

5

20

- 1. A pharmaceutical aerosol formulation comprising
- (a) 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoro-n-propane or mixtures thereof as propellant;
- (b) 1,1,2,2,3-pentafluoropropane as ∞-propellant; and
- (c) particulate medicament.
- 2. A formulation according to claim 1 wherein the ratio of propellant:copropellant is about 30:70 to about 95:5 by weight.
 - 3. A formulation according to claim 2 wherein the ratio of propellant:co-propellant is about 50:50 to about 80:20 by weight.
- 15 4. A formulation according to any one of claims 1 to 3 wherein the propellant comprises 1,1,1,2-tetrafluoroethane.
 - 5. A formulation according to any one of claims 1 to 3 wherein the propellant comprises 1,1,1,2,3,3,3-heptafluoro-n-propane.
 - 6. A formulation according to any one of claims 1 to 5 wherein the medicament is an anti-allergic, a bronchodilator or an anti-inflammatory steroid.
 - 7. A formulation according to any one of claims 1 to 6 wherein the medicament is salmeterol xinafoate.
 - 8. A formulation according to any one of claims 1 to 6 wherein the medicament is salbutamol sulphate.
- 30 9. A formulation according to any one of claims 1 to 6 wherein the medicament is fluticasone propionate.

- 10. A formulation according to any one of claims 1 to 6 wherein the medicament is becomethasone dipropionate of a physiolgically acceptable solvate thereof.
- 11. A formulation according to any one of claims 1 to 6 wherein the medicament is formoterol, cromoglycate, terbutaline, reproterol or (-)-4-amino-3,5-dichloro-α-[[[6-[2-(2-pyridinyl)ethoxy]hexyl]amino]methyl]benzemethanol budesonide, triamcinolone acetonide or a physiologically acceptable salt or solvate thereof.
 - 12. A formulation according to any one of claims 1 to 11 wherein the medicament is present in an amount of 0.005 to 10% w/w relative to the total weight of the formulation.
- 13. A formulation according to claim 12 wherein the medicament is present in an amount of 0.01 to 1% w/w relative to the total weight of the formulation.
 - 14. A formulation according to any one of claims 1 to 13 which contains two or more particulate medicaments.
- 20
 15. A formulation according to claim 14 which contains salbutamol or salmeterol or a physiologically acceptable salt thereof in combination with an anti-inflammatory steroid or an anti-allergic.
- 16. A formulation according to claim 15 which contains salmeterol or salbutamol or a physiologically acceptable salt thereof in combination with fluticasone propionate or beclomethasone dipropionate or a physiologically acceptable solvate thereof.
- 17. A formulation according to any one of claims 1 to 16 comprising an adjuvant having a higher polarity and/or a boiling point than the propellant.
 - 18. A formulation according to claim 17 wherein the adjuvant having a higher polarity than the propellant is present in an amount of 0.05 to 5% w/w based upon the propellant and co-propellant.

- 19. A formulation according to any one of claims 1 to 18 comprising a surfactant.
- 5 20. A canister suitable for delivering a pharmaceutical aerosol formulation which comprises a container capable of withstanding the vapour pressure of the propellant used which container is closed with a metering valve and contains a pharmaceutical aerosol formulation which comprises (a) 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoro-n-propane or mixtures thereof as propellant, (b) 1,1,2,2,3-pentafluoropropane as co-propellant, and (c) a particulate medicament.
 - 21. A canister according to claim 20 wherein the container is a metal can.
- 15 22. A canistrer according to claim 21 wherein the container is an aluminium can.
 - 23. A canister according to claim 21 or 22 wherein the container is plastics-coated.
 - 24. A metered dose inhaler which comprises a canister according to any one of claims 20 to 24 fitted into a suitable channelling device.
- 25. A method of treating respiratory disorders which comprises administration by inhalation of an effective amount of a pharmaceutical aerosol formulation according to any one of claims 1 to 18.

Int Jonal Application No

			PCT/EP 95/04824
A. CLAS	SIFICATION OF SUBJECT MATTER A61K9/00 A61K9/12		
According	to International Patent Classification (IPC) or to both national	I classification and IDC	
B. FIELD	DS SEARCHED	,	
Minimum IPC 6	documentation searched (damification system followed by da A61K	mification symbols)	
			·
Document	ation scarched other than minimum documentation to the exten	it that such documents are includ	ed in the fields searched
Electronic	data base consulted during the international search (name of di	the base and sub-re-	
	Colora Colo	· · · · · · · · · · · · · · · · · · ·	rich (cima used)
	·,		
C. DOCUM	MENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of	the relevant passages	Relevant to claim No.
Υ .	WO.A.93 02150 (DU PONT) 4 Febr see page 13, column 6, line 8 table 1	uary 1993 - line 12:	1-25
Y	US,A,5 314 926 (ROBIN MARK L ET AL) 24 May 1994		1-25
	see column 3, line 1 - line 61		
Y	WO,A,94 03153 (GLAXO GROUP LTD ANTHONY JAMES (GB); NEALE PHIL (GB)) 17 February 1994 see claims 1-13	;TAYLOR IP JOHN	1-25
γ	WO,A,93 11745 (GLAXO GROUP LTD) 1993 see claims 1-16) 24 June	1-25
	•••	-/	
[
X Furth	er documents are listed in the continuation of box C.	X Patent (amily memb	bers are listed in annex.
Special care	gones of cited documents:	T later document authlished	d after the international filing date
COURGEL	nt defining the general state of the art which is not red to be of particular relevance ocument but published on or after the international	or priority data and not	or are the members application but in conflict with the application but principle or theory underlying the
filing da L' documen which is	ite . It which may throw doubts on priority claim(s) or cited to exabilish the publication date of another	cannot be considered no involve an inventive ste	relevance; the claimed invention over or cannot be considered to p when the document in taken alone relevance; the claimed invention
O° documen other me		cannot be considered to document is combined to ments, such combination	n being obvious to a person skilled
later than	t published prior to the international filing date but n the priority date clarmed	in the art, "A" document member of th	e same patent family
ate of the ac	tual completion of the international search		nternational search report
11	March 1996		02.04.96
ame and ma	iling address of the ISA European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk	Authorized officer	
	Tcl. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Foerster,	W

Form PCT/ISA/210 (second sheet) (July 1992)

Int Jonal Application No PCT/EP 95/04824

Category *	non) DOCUMENTS CONSIDERED TO BE RELEVANT Clunon of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,93 11743 (GLAXO GROUP LTD) 24 June	1-25
	1993 see claims 1-22	
A	WO,A,93 18746 (ASTA MEDICA AG) 30 September 1993 see claims 1-12	1-25
١	WO.A.91 14422 (MINNESOTA MINING & MFG) 3 October 1991 cited in the application see claims 1-9	1-25
	•••••	
•		
	•	
Ì		
ł		
ŀ		
	•	
- [
.]		
		·
1		

Information on patent family members

Int. Jonal Application No PCT/EP 95/04824

Patent document cited in search report	Publication date	Pater mer	Patent family member(s)		
WO-A-9302150	04-02-93	AU-B- 2343692		23-02-93	
		CN-A-	1068841	10-02-93	
		DE-D-	69204290	28-09-95	
•		EP-A-	0595937	11-05-94	
		EP-A-	0661365	05-07-95	
US-A-5314926	24-05-94	US-A-	5278196	11-01-94	
•	•	AU-B-	6589794	11-10-94	
		EP-A-	0690887	10-01-96	
		WO-A-	9421718	29-09-94	
		ZA-A-	9401859	17-10-94	
√0-A-9403153	17-02-94	AU-B-	4705093	03-03-94	
		CA-A-	2141039	17-02-94	
•	•	CN-A-	1088436	29-06-94	
•		EP-A-	0658101	21-06-95	
		JP-T-	7509475	19-10-95	
		ZA-A-	9305477	23-02-94	
0-A-9311745	24-06-93	AT-T-	128350	15-10-95	
		AU-B-	663986	26-10-95	
		AU-B-	3085292	19-07-93	
		CA-A-	2125665	24-05-93	
	•	DE-D-	69205177	02-11-95	
		EP-A-	0616525	28-09-94	
•	•	JP-T-	7501811	23-02-95	
•		NZ-A-	246046	21-12-95	
		ZA-A-	9209618	09-08-93	
		AP-A-	402	22-08-95	
		AU-B-	663904	26-10-95	
		AU-B-	3085092	19-07-93	
		AU-B-	663905	26-10-95	
•		AU-B-	3085192	19-07-93	
	•	BG-A-	98803	28-02-95	
	• •	CA-A-	2125666 ⁻	24-06-93	
		CA-A-	2125667	24-06-93	
•		CN-A-	1075078	11-08-93	
	•	CN-A-	1075079	11-08-93	
		CZ-A-	9401430	15-03-95	
. •	•	WO-A-	9311743	24-06-93	

Information on patent family members

In. ional Application No PCT/EP 95/04824

Patent document cited in search report	Publication date		t family iber(s)	Publication date	
WO-A-9311745		WO-A-	9311744	24-06-93	
NUTA : JUAAT TU		EP-A-	0616523	28-09-94	
		EP-A-	0616524	28-09-94	
		HU-A-	67534	28-04-95	
		JP-T-	7502033	02-03-95	
	·	JP-T-	7502034	02-03-95	
		NO-A-	942185	10-06-94	
		NZ-A-	246044	26-01-96	
		OA-A-	9926	15-09-94	
		SK-A-	67494	08-03-95	
WO-A-9311743	24-06-93	AP-A-	482	22-08-95	
•		AU-B-	663904	26-10-95	
:		AU-B-	3085 092	19-07-93	
1		BG-A-	98803	28-82-95	
•	•	CA-A-	2125667	24-06-93	
· ·		CZ-A-	9401430	15-83-95	
		EP-A-	9616523	28-09-94	
		HU-A-	67534	28-04-95	
		JP-T-	7502033	02-03-95	
		NO-A-	942185	10-06-94	
		NZ-A-	246044	26-01-96	
		OA-A-	9926	15-09-94	
		SK-A-	67494	08-03-95	
	•	ZA-A-	9209617	22-03-94	
		AU-B-	663905	26-10-95	
•		AU-B-	3085192	19-07-93	
		CA-A-	2125666	24-06-93	
	•	WO-A-	9311744	24-06-93	
		EP-A-	0616524	28-09-94	
		JP-T-	7502034	02-03-95	
*		AT-T-	128350	15-10-95	
		AU-B-	663906	26-10-95	
		AU-B-	3085292	19-07-93	
	. ,	CA-A-	2125665	24-06-93	
		CN-A-	1075078	11-08-93	
		CN-A-	1075079	11-08-93	
		DE-D-	69205177	02-11-95	
		WO-A-	9311745	24-06-93	
		EP-A-	0616525	28-09-94	
				. •	
				,	
		•			
,					
				a .	
				•	
				•	

Information on patent family members

In. Jonal Application No PCT/EP 95/04824

Patent document cited in search report	Publication date	Patent family member(s)		Publication date	
WO-A-9311743		JP-T-	7501811	23-02-95	
		NZ-A-	246046	21-12-95	
WO-A-9318746	30-09-93	DE-A-	4230876	23-09-93	
		AU-B-	3745993	21-10-93	
		CA-A-	2129855	18-09-93	
		EP-A-	0630229	28-12-94	
		FI-A-	944257	14-69-94	
		HU-A-	68223	28-06-95	
		JP-T-	7508506	21-09-95	
		NO-A-	943305	07-09-94	
	•	SK-A-	385892	12-04-95	
		US-A-	5415853	16-05-95	
		ZA-A-	9301907	06-10-93	
WO-A-9114422	03-10-91	AU-B-	654813	24-11-94	
	· · · · · · · · · · · · · · · · · · ·	AU-B-	7668691	21-10-91	
		CA-A-	2077354	24-69-91	
•		DE-D-	69109284	01-06-95	
•	•	DE-T-	69109284	24-08-95	
		EP-A-	0526481	10-02-93	
•		EP-A-	0636362	01-02-95	
		ES-T-	2071306	16-06-95	
		US-A-	5118494	02-06-92	